

BIBLIOGRAFIA CAPITOLO HIGH TECH

RADIOFREQUENCY

1. Alvaro P, Alberto O, Matia F, Beatrice Marina P. An innovative temperature-controlling handpiece for face and body skin laxity and tightening treatment with radiofrequency. *Skin Res Technol*. 2023 Jun;29(6):e13385. doi: 10.1111/srt.13385. PMID: 37357641; PMCID: PMC10246699.
2. Levenberg A. Clinical experience with a TriPollar radiofrequency system for facial and body aesthetic treatments. *Eur J Dermatol*. 2010; 20(5): 615-619. <https://doi.org/10.1684/EJD.2010.1042>
3. Duncan DI (2012) Improving outcomes in upper arm liposuction: adding radiofrequency-assisted liposuction to induce skin contraction. *Aesthet Surg J* 32(1):84–95. <https://doi.org/10.1177/1090820X11429549>
4. Katz B (2023) Concomitant use of radiofrequency and high intensity focused electromagnetic field energies for full-body remodeling: MRI evidence-based preface trial. *J Cosmet Dermatol* 22:193–199. <https://doi.org/10.1111/jocd.14736>
5. Park JH, Kim JI, Park HJ, Kim WS. Evaluation of safety and efficacy of noninvasive radiofrequency technology for submental rejuvenation. *Lasers Med Sci*. 2016 Nov;31(8):1599-1605. doi: 10.1007/s10103-016-2023-7. Epub 2016 Jul 12. PMID: 27402002.
6. Locketz GD, Bloom JD. Percutaneous Radiofrequency Technologies for the Lower Face and Neck. *Facial Plast Surg Clin North Am*. 2019 Aug;27(3):305-320. doi: 10.1016/j.fsc.2019.03.003. Epub 2019 May 16. PMID: 31280845.
7. Azuelos A, SidAhmed-Mezi M, La Padula S, Aboud C, Meningaud JP, Hersant B. High-Intensity Focused Ultrasound: A Satisfactory Noninvasive Procedure for Neck Rejuvenation. *Aesthet Surg J*. 2019 Jul 12;39(8):NP343-NP351. doi: 10.1093/asj/sjz093. PMID: 30923813.
8. Tettamanzi M, Pili N, Rodio M, Serra PL, Trignano C, Rubino C, Trignano E. Use of Radiofrequency-Assisted Liposuction (BodyTite) for Upper Arms Lifting. *Aesthetic Plast Surg*. 2023 Dec;47(6):2495-2501. doi: 10.1007/s00266-023-03452-6. Epub 2023 Jun 14. PMID: 37314467; PMCID: PMC10784384.

HIGH TECH IN FACIAL PLASTIC SURGERY

1. Feldman LS, Fuchshuber PR, Jones DB. The SAGES manual on the fundamental use of surgical energy (FUSE). Berlin/Heidelberg (Germany): Springer Science+Business Media, LLC; 2012.
2. Anderson RR, Parrish JA. Selective photothermolysis: precise microsurgery by selective absorption of pulsed radiation. *Science* 1983;220(4596):524–7.
3. Holcomb JD, Rousso KK, Rousso DE. Nitrogen plasma skin regeneration and aesthetic facial surgery. Multicenter Evaluation of Concurrent Treatment. *Arch Facial Plast Surg* 2009;11(3): 184–93.
4. Kilmer S, Shah SN, Fitzpatrick G, et al. A pilot study on the use of a plasma skin regeneration device (Portrait!PSR3) in full facial rejuvenation procedures. *Lasers Med Sci* 2007;22(2):101–9.

5. Gentile RD. Cool atmospheric plasma (J-plasma) and new options for facial contouring and skin rejuvenation of the heavy face and neck. *Facial Plast Surg* 2018;34(1):66–74.
6. Gentile RD. Renuvion/J-plasma for subdermal skin tightening facial contouring and skin rejuvenation of the face and neck. *Facial Plast Surg Clin North Am* 2019;27(3):273–90.
7. Nistico SP, Chiricozzi A, Tamburi F, et al. Lasers and energy devices for the skin: conventional and unconventional use. *Biomed Res Int* 2016;2016: 9031091. Accessed June 1, 2019.
8. Gentile RD. Smartlifting-A technological innovation for facial rejuvenation. *Lasers Surg Med* 2009; 41(Supplement 21).
9. Gentile RD. SmartLifting fiber laser-assisted facial rejuvenation techniques. *Facial Plast Surg Clin North Am* 2011;19(02):371–87.
10. Gentile RD. Subcutaneous fiber laser and energy- based techniques for facial rejuvenation. In: Truswell WH, editor. *Lasers and light, peels and abrasions*. New York: Thieme Publishers; 2016. p. 47–74.
11. Gentile RD. Laser-assisted neck-lift: high-tech contouring and tightening. *Facial Plast Surg* 2011; 27(4):331–45.

LASERS AND SCARS

1. Nguyen TA, Feldstein SI, Shumaker PR, Krakowski AC. A review of scar assessment scales. *Semin Cutan Med Surg* 2015;34(1):28–36.
2. Dalkey N. The Delphi Method: An experimental study of group opinion. *Futures* 1969;1:408–426.
3. Wood F. An approach to scar mitigation. In: Krakowski A, Shumaker P, editors. *The Scar Book*. Philadelphia, PA: Wolters Kluwer; 2017.
4. Tredget E, Ding J. The cellular and molecular basis of scarring: The paradigm of hypertrophic scarring after thermal injury. In: Krakowski A, Shumaker P, editors. *The Scar Book*. Philadelphia, PA: Wolters Kluwer; 2017.
5. Carlavan I, Bertino B, Rivier M, et al. Atrophic scar formation in patients with acne involves long-acting immune responses with plasma cells and alteration of sebaceous glands. *Br J Dermatol* 2018;179(4):906–917.
6. Ansari F, Sadeghi-Ghyassi F, Yaaghoobian B. The clinical effectiveness and cost-effectiveness of fractional CO₂ laser in acne scars and skin rejuvenation: A meta-analysis and economic evaluation. *J Cosmet Laser Ther* 2018;20(4):248–251.
7. Xu Y, Deng Y. Ablative fractional CO₂ laser for facial atrophic acne scars. *Facial Plast Surg* 2018;34(2):205–219.
8. Dai R, Xie H, Hua W, Li XH, Li L. The efficacy and safety of the fractional radiofrequency technique for the treatment of atrophic acne scar in Asians: A meta-analysis. *J Cosmet Laser Ther* 2017;19(6):337–344.
9. Ozog DM, Liu A, Chaffins ML, et al. Evaluation of clinical results, histological architecture, and collagen expression following treatment of mature burn scars with a fractional carbon dioxide laser. *JAMA Dermatol* 2013;149(1):50–57.
10. Connolly KL, Chaffins M, Ozog D. Vascular patterns in mature hypertrophic burn scars treated with fractional CO₂ laser. *Lasers Surg Med* 2014;46(8):597–600.
11. Qu L, Liu A, Zhou L, et al. Clinical and molecular effects on mature burn scars after treatment with a fractional CO₂ laser. *Lasers Surg Med* 2012;44(7):517–524.

12. Onur Erol O, Atabay K. The treatment of burn scar hypopigmentation and surface irregularity by dermabrasion and thin skin grafting. *Plast Reconstr Surg* 1990;85(5):754–758.
13. Waibel JS, Wulkan AJ, Shumaker PR. Treatment of hypertrophic scars using laser and laser assisted corticosteroid delivery. *Lasers Surg Med* 2013;45(3):135–140.
14. Waibel J, Wulkan AJ, Lupo M, Beer K, Anderson RR. Treatment of burn scars with the 1,550 nm nonablative fractional Erbium Laser. *Lasers Surg Med* 2012;44(6):441–446.
15. Bowes LE, Nouri K, Berman B, et al. Treatment of pigmented hypertrophic scars with the 585 nm pulsed dye laser and the 532 nm frequency-doubled Nd:YAG laser in the Q-switched and variable pulse modes: a comparative study. *Dermatol Surg* 2002;28(8):714–719.
16. Hultman CS, Friedstat JS, Edkins RE. Efficacy of intense pulsed light for the treatment of burn scar dyschromias: a pilot study to assess patient satisfaction, safety, and willingness to pay. *Ann Plast Surg* 2015;74(Suppl 4):S204–S208.
17. Massaki AB, Fabi SG, Fitzpatrick R. Repigmentation of hypopigmented scars using an erbium-doped 1,550-nm fractionated laser and topical bimatoprost. *Dermatol Surg* 2012;38(7 Pt 1):995–1001.
18. Goodman GJ. An automated autologous cell transplantation method for the treatment of hypopigmented scarring. *Dermatol Surg* 2008;34(4):578–581.
19. Jin R, Huang X, Li H, et al. Laser therapy for prevention and treatment of pathologic excessive scars. *Plast Reconstr Surg* 2013;132(6):1747–1758.
20. Al-Mohamady Al-S, Ibrahim SM, Muhammad MM. Pulsed dye laser versus long-pulsed Nd:YAG laser in the treatment of hypertrophic scars and keloid: A comparative randomized split-scar trial. *J Cosmet Laser Ther* 2016;18(4):208–212.
21. Keaney TC, Tanzi E, Alster T. Comparison of 532 nm potassium titanyl phosphate laser and 595 nm pulsed dye laser in the treatment of erythematous surgical scars: A randomized, controlled, open-label study. *Dermatol Surg* 2016;42(1):70–76.